

Contents lists available at [ScienceDirect](http://ScienceDirect.com)

Journal of Arrhythmia

journal homepage: www.elsevier.com/locate/joa

Review

Ventriculoventricular delay optimization of a cardiac resynchronization device



Koichi Sagara, MD*

Department of Cardiovascular Medicine, The Cardiovascular Institute, 3-2-19 Nishiazabu, Minato-ku, Tokyo, Japan

ARTICLE INFO

Article history:

Received 17 January 2014

Accepted 24 March 2014

Available online 12 June 2014

Keywords:

Cardiac resynchronization therapy

VV delay

Interventricular interval

ABSTRACT

Cardiac resynchronization therapy (CRT) has become a standard option for patients with severe low cardiac function and mild to severe heart failure. However, its potential has not been maximized to date, as the optimal atrioventricular delay, ventriculoventricular (VV) delay, and tachy therapy settings remain unknown. Here, data from various studies have been used to estimate several CRT settings.

Three search words—interventricular interval, VV delay, and interventricular delay with cardiac resynchronization therapy—were entered into PubMed. The methods used to optimize VV delay included ultrasonography, radioisotope diagnosis, scintigraphy, electrocardiography, Swan–Ganz catheterization, and thoracic impedance. Their populations and results were analyzed to identify convincing rules. Methods for VV delay optimization in the literature can be categorized into four patterns. Time and cost were high in several categories. Most studies concluded that their method was effective but no small amount of papers denied individual detailed optimization. There were some population biases in most papers. Individual optimization had a major impact in patients with ischemic heart disease but no significant impact in patients with non-ischemic heart diseases.

In summary, CRT is an established therapy, but a well-controlled study is required to find conclusive methods for VV delay optimization.

© 2014 Japanese Heart Rhythm Society. Published by Elsevier B.V. All rights reserved.

Contents

1. Introduction	413
1.1. Cardiac resynchronization therapy (CRT)	413
1.2. Categorization of VV delay optimization methods	414
2. Methods and results	414
3. Conclusion	414
Conflict of interest	415
Acknowledgment	415
References	415

1. Introduction

Heart failure is a very common life-threatening disease that incurs a high cost. Therapy for heart failure is important because it not only improves patients' quality of life and prognosis but may also prevent subsequent hospitalization and reduce social medical costs. Optimal therapy for heart failure is needed from both the social and individual perspectives.

1.1. Cardiac resynchronization therapy (CRT)

CRT is firmly established but has many areas in need of improvement [1–12]. Particularly, the prognostic value to predict who responds to CRT is not well established [13–19]. Only QRS duration has a confirmed value, the sensitivity of which is insufficient, however [20–22]. The cost of CRT, although more economically reasonable than other therapies, is high; hence, a more effective method to predict responders needs to be established.

In addition, the ideal CRT optimization method is unknown [23]. There are several effective methods, but one has yet to be

* Tel.: +81 3 3408 2151.

E-mail address: k-sagara@cvi.or.jp

identified as the best [24–28]. Especially regarding VV delay, there are many conflicting methods and theories [29–31].

1.2. Categorization of VV delay optimization methods

Methods to optimize VV delay remain competitive; therefore, four categories were created in which to summarize them. The first category focuses on wall motion and dyssynchrony using ultrasonography, scintigraphy, and other techniques [2,3,32–48]. The second category focuses on conduction delays and disorders using electrocardiography and a calculation of lead-to-lead conduction delay time (time between right ventricular pacing and left ventricular sensing and between left ventricular pacing and right ventricular sensing), an electroanatomic mapping system, and other techniques [22,49–63]. The third category considers cardiac output estimated by Swan–Ganz catheterization, thoracic impedance, ultrasonography, and other techniques [55,57,58,64–70]. The fourth category eliminates individual optimization and uses a standard value because these methods acknowledge no major differences in VV delay settings [3,35,71–82].

Since CRT is pacing therapy, it may correct conduction delays but does not improve muscle function. Accordingly, the second category was most appropriate, but most major methods use ultrasonography and a majority of papers in this area claim method efficacy. This may be why we did not want to improve conduction delays; rather, our aim was to relieve heart failure symptoms by increasing cardiac output. The third category matched this aim but in many cases required extensive time and cost. And we decide optimal settings at rest, which may not be optimal during activity time. We do not know whether optimal settings at rest improve a patient's activity and prognosis or not. The fourth category solves this problem: if optimization is not required, it may be the best option.

The first step of this systematic review was determining the need for optimization, followed by identifying the best method to optimize the VV delay.

2. Methods and results

A literature search using the PubMed/Medline databases and consensus documents was performed. Searches using “interventricular interval” found 43 studies, “VV delay” found 59 studies, and “interventricular delay” found 200 studies of CRT.

In each study, optimization need, associated methods, optimization modality, and populations were counted. Whenever possible, optimization need as well as method and modality superiority was decided objectively (e.g., the measurement of velocity time integral was not better than the measurement of QRS duration).

A total of 33 studies confirmed the need for optimization and clearly described the superiority of a detailed optimization. These studies used the velocity time integral measured by ultrasonography, the spectrum of wall motion, dyssynchronicity, nuclear imaging, electrocardiography, and other methods. Their superiority was proven by both the instant outcome and the clinical results.

Five studies denied the need for optimization. The nominal settings, including a fixed VV delay ($LV > RV$ of 40 ms or $LV = RV$), met the clinical purpose. They tended not to measure small details, perhaps since optimal settings change in different situations and small details are of little consequence. If the clinical output is unchanged, there is no need to optimize the VV delay.

Seven studies partially affirmed the need for optimization by removing a detailed check and programing the calculations using pre-populated generator programing (e.g., Quick Opt, St. Jude Medical, Inc.)

The use of ultrasonographic cardiography, the most common modality, for optimization was recommended by 25 studies. The

estimation of cardiac output using velocity time integral measurements was most common. Conduction disorders and ventricular muscle dyssynchrony were also improved by spectral measurements, septal wall motion delays, and other techniques.

Ten studies emphasized the importance of conduction delays and stated that electrocardiography was a superior method. In fact, the smallest QRS duration was determined using electrocardiography. There is clear evidence that a wide QRS complex is a prognostic factor of CRT, but it is not clear whether a narrow QRS pacing is preferable.

Seven studies stated that the use of pre-populated programing is both necessary and sufficient. Clinically speaking, it met the demand, but it was unable to obtain the best settings. Therefore, its use might be sufficient for clinical situations.

The ability to obtain a detailed understanding of cardiac function using Swan–Ganz catheterization, thoracic impedance, and other methods has been proven, but few studies have proven their superiority; rather, they stated that cardiac output was changed by VV delay but did not determine its influence on clinical output [65,68].

The populations of these studies had various biases. Despite relatively slight differences in age, sex, and QRS duration, in some studies, > 80% of patients had ischemic heart disease, whereas in others, > 70% of patients had non-ischemic heart disease.

The majority of these studies concluded that their particular method was effective, but most denied that their method was individual, detailed, and had universal optimization. In most studies, the population tended to be weighted in one direction. Individual optimization had some impact in patients with ischemic heart disease but no major impact in patients with non-ischemic heart disease.

In the present study, the effect of population bias was particularly considered. A number of papers that argued that a detailed VV delay optimization improves heart failure contained a large proportion of patients with ischemic heart disease. On the other hand, many papers that claimed that detailed optimization was not important contained a large proportion of patients with non-ischemic heart disease. Hence, it is possible that a detailed VV delay optimization was effective in patients with ischemic heart disease and not needed in patients with non-ischemic heart disease.

CRT is a therapy for conduction disorders. Damaged cardiomyocytes are localized in ischemic heart disease; therefore, in a way, the presence of a conduction disorder is localized and varies among the cases. On the other hand, conduction disorders were not always unequal in patients with non-ischemic heart disease. Localized and inhomogeneous conduction disorders require individualized conduction therapy with CRT, whereas diffuse conduction disorders require only standard settings.

3. Conclusion

CRT is a firmly established therapy, but we do not yet know how to maximize its potential at minimum cost. CRT is a costly therapy; hence, its potential should be fully utilized. We should also determine routine CRT settings because its use is becoming more common and an increasing number of CRT devices will be implanted in the future.

Here, VV delay settings, which are very common but controversial, were particularly noted. Upon examining the study findings, a therapy for conduction disorder indicated from the conduction perspective may be decided by conduction delay dispersion.

Conduction disorders were localized and inhomogeneous in patients with ischemic heart disease, in whom CRT efficacy was slightly low; therefore, detailed optimization using echocardiography,

radioisotopes, Swan–Ganz catheterization, and other methods should be carefully performed as needed [83,84].

Dilated cardiomyopathy (DCM) has a relatively homogeneous diffuse conduction order. Clinically simultaneous biventricular pacing was the most common setting for DCM patients. In the future, we can determine the indication for optimization according to specific diseases or heart conditions.

Here, methods for interventricular delay optimization were categorized into four groups. The first category emphasized wall motion and dyssynchrony and used ultrasonography. If dyssynchrony was judged to be high, VV delay optimization using ultrasonography was needed. The second focused heavily on conduction delays and disorders using electrocardiography. If the QRS duration was very wide or had a left bundle branch block pattern, VV delay optimization was needed. Signal-averaged electrocardiography was also useful, but the meaning of its results was uncertain [85]. The third category placed great weight on cardiac output. In patients with very low cardiac output, conduction disorders tended to be high and diffuse and not require optimization. Similarly, in the fourth category, heart disease in which a diffuse conduction disorder was implied did not require optimization. In any case, the need for VV delay optimization might be judged according to a specific disease or conduction disorder pattern; hence, the most effective and least costly settings should be identified.

Conflict of interest

There is no conflict of interest.

Acknowledgment

No special instruction.

References

- [1] Achilli A, Sassara M, Ficili S, et al. Long-term effectiveness of cardiac resynchronization therapy in patients with refractory heart failure and narrow QRS. *J Am Coll Cardiol* 2003;42:2117–24.
- [2] Chan KL, Tang AS, Achilli A, et al. Functional and echocardiographic improvement following multisite biventricular pacing for congestive heart failure. *Can J Cardiol* 2003;19:387–90.
- [3] Wang D, Yu H, Yun T, et al. Long-term clinical effects of programmer-guided atrioventricular and interventricular delay optimization: intracardiac electrography versus echocardiography for cardiac resynchronization therapy in patients with heart failure. *J Internat Med Res* 2013;41:115–22.
- [4] Wang A, Cabrera SE, Havalad V, et al. Effects of biventricular pacing on left heart twist and strain in a porcine model of right heart failure. *J Surg Res* 2013;185:645–52.
- [5] Spotnitz ME, Richmond ME, Quinn TA, et al. Relation of QRS shortening to cardiac output during temporary resynchronization therapy after cardiac surgery. *ASAIO J* 2010;56:434–40.
- [6] Wang DY, Richmond ME, Quinn TA, et al. Optimized temporary biventricular pacing acutely improves intraoperative cardiac output after weaning from cardiopulmonary bypass: a substudy of a randomized clinical trial. *J Thorac Cardiovasc Surg* 2011;141:1002–8, 8.e1.
- [7] Spotnitz HM, Cabrera SE, Wang DY, et al. Primary endpoints of the biventricular pacing after cardiac surgery trial. *Ann Thorac Surg* 2013;96:808–15.
- [8] Celebi O, Knaus T, Blaschke F, et al. Extraordinarily favorable left ventricular reverse remodeling through long-term cardiac resynchronization: super-response to cardiac resynchronization. *Pacing Clin Electrophysiol* 2012;35:870–6.
- [9] Dreger H, Maethner K, Bodke H, et al. Pacing-induced cardiomyopathy in patients with right ventricular stimulation for > 15 years. *Europace* 2012;14:238–42.
- [10] Luo C, Li Y, Liu D, et al. The association of brachial flow-mediated dilation and high-sensitivity C-reactive protein levels with Duke treadmill score in patients with suspected microvascular angina. *Exp Clin Cardiol* 2012;17:197–201.
- [11] Sitges M, Vidal B, Delgado V, et al. Long-term effect of cardiac resynchronization therapy on functional mitral valve regurgitation. *Am J Cardiol* 2009;104:383–8.
- [12] Ghio S, Freemantle N, Scelsi L, et al. Long-term left ventricular reverse remodelling with cardiac resynchronization therapy: results from the CARE-HF trial. *Eur J Heart Fail* 2009;11:480–8.
- [13] Aaronaes M, Aakhus S, Aass H, et al. Assessment of response criteria to cardiac resynchronization therapy (CRT) and prediction of response. *Scand Cardiovasc J* 2010;44:337–45.
- [14] van Bommel RJ, Borleffs CJ, Ypenburg C, et al. Morbidity and mortality in heart failure patients treated with cardiac resynchronization therapy: influence of pre-implantation characteristics on long-term outcome. *Eur Heart J* 2010;31:2783–90.
- [15] van Bommel RJ, Ypenburg C, Borleffs CJ, et al. Value of tissue Doppler echocardiography in predicting response to cardiac resynchronization therapy in patients with heart failure. *Am J Cardiol* 2010;105:1153–8.
- [16] van Bommel RJ, Tanaka H, Delgado V, et al. Association of intraventricular mechanical dyssynchrony with response to cardiac resynchronization therapy in heart failure patients with a narrow QRS complex. *Eur Heart J* 2010;31:3054–62.
- [17] Zucchelli G, Soldati E, Di Cori A, et al. Role of intraoperative electrical parameters in predicting reverse remodelling after cardiac resynchronization therapy and correlation with interventricular mechanical dyssynchrony. *Europace* 2010;12:1453–9.
- [18] Bordachar P, Lafitte S, Reant P, et al. Low value of simple echocardiographic indices of ventricular dyssynchrony in predicting the response to cardiac resynchronization therapy. *Eur J Heart Fail* 2010;12:588–92.
- [19] Wiesbauer F, Baytaroglu C, Azar D, et al. Echo Doppler parameters predict response to cardiac resynchronization therapy. *Eur J Clin Invest* 2009;39:1–10.
- [20] Brenyo A, Rao M, Barsheshet A, et al. QRS axis and the benefit of cardiac resynchronization therapy in patients with mildly symptomatic heart failure enrolled in MADIT-CRT. *J Cardiovasc Electro* 2013;24:442–8.
- [21] Chapurnykh AV, Mochalova OV, Solov'eva NV, et al. Cardiac resynchronization therapy in patients with narrow QRS. *Kardiologiya* 2011;51:82–8.
- [22] Tamborero D, Vidal B, Tolosana JM, et al. Electrocardiographic versus echocardiographic optimization of the interventricular pacing delay in patients undergoing cardiac resynchronization therapy. *J Cardiovasc Electrophysiol* 2011;22:1129–34.
- [23] Qamruddin S, Rafie R, Naqvi TZ. Discordant left and right ventricular optimal atrioventricular and interventricular delays during biventricular pacemaker optimization. *Echocardiography* 2013;30:751–8.
- [24] Risum N, Sogaard P, Hansen TF, et al. Comparison of dyssynchrony parameters for VV-optimization in CRT patients. *Pacing Clin Electrophysiol* 2013;36:1382–90.
- [25] Birnie D, Lemke B, Aonuma K, et al. Clinical outcomes with synchronized left ventricular pacing: analysis of the adaptive CRT trial. *Heart Rhythm* 2013;10:1368–74.
- [26] Bertini M, Valzania C, Biffi M, et al. Interventricular delay optimization: a comparison among three different echocardiographic methods. *Echocardiography* 2010;27:38–43.
- [27] Ypenburg C, van Bommel RJ, Delgado V, et al. Optimal left ventricular lead position predicts reverse remodeling and survival after cardiac resynchronization therapy. *J Am Coll Cardiol* 2008;52:1402–9.
- [28] Ypenburg C, Van De Veire N, Westenberg JJ, et al. Noninvasive imaging in cardiac resynchronization therapy—Part 2: Follow-up and optimization of settings. *Pacing Clin Electrophysiol* 2008;31:1628–39.
- [29] Abraham WT, Leon AR, Sutton St John, et al. Randomized controlled trial comparing simultaneous versus optimized sequential interventricular stimulation during cardiac resynchronization therapy. *Am Heart J* 2012;164:735–41.
- [30] Abraham WT, Smith SA. Devices in the management of advanced, chronic heart failure. *Nat Rev Cardiol* 2013;10:98–110.
- [31] Houthuizen P, Bracke FA, van Gelder BM. Atrioventricular and interventricular delay optimization in cardiac resynchronization therapy: physiological principles and overview of available methods. *Heart Fail Rev* 2011;16:263–76.
- [32] Harel F, Finnerty V, Gregoire J, et al. Effects of dobutamine stress on cardiac contraction synchronism in a canine model. *Physiol Meas* 2013;34:1387–97.
- [33] Dong YX, Oh JK, Yang YZ, et al. Left ventricular 12 segmental strain imaging predicts response to cardiac resynchronization therapy. *Chin Med J* 2013;126:2620–4.
- [34] Praus R, Haman L, Tauchman M, et al. Echocardiographic changes after cardiac resynchronization therapy. *Kardiol Pol* 2012;70:1250–7.
- [35] Pabari PA, Willson K, Stegmann B, et al. When is an optimization not an optimization? Evaluation of clinical implications of information content (signal-to-noise ratio) in optimization of cardiac resynchronization therapy, and how to measure and maximize it. *Heart Fail Rev* 2011;16:277–90.
- [36] Kyriacou A, Pabari PA, Mayet J, et al. Cardiac resynchronization therapy and AV optimization increase myocardial oxygen consumption, but increase cardiac function more than proportionally. *Int J Cardiol* 2014;171:144–52.
- [37] Kyriacou A, Li Kam Wa ME, Pabari PA, et al. A systematic approach to designing reliable VV optimization methodology: assessment of internal validity of echocardiographic, electrocardiographic and haemodynamic optimization of cardiac resynchronization therapy. *Int J Cardiol* 2013;167:954–64.
- [38] Tatsumi K, Tanaka H, Tsuji T, et al. Strain dyssynchrony index determined by three-dimensional speckle area tracking can predict response to cardiac resynchronization therapy. *Cardiovasc Ultrasound* 2011;9:11.
- [39] Tatsumi K, Tanaka H, Yamawaki K, et al. Utility of comprehensive assessment of strain dyssynchrony index by speckle tracking imaging for predicting response to cardiac resynchronization therapy. *Am J Cardiol* 2011;107:439–46.
- [40] Naqvi TZ. Echocardiography-guided biventricular pacemaker optimization. *JACC Cardiovasc Imaging* 2010;3:1168–80.

- [41] Gorcsan 3rd J, Oyenuga O, Habib PJ, et al. Relationship of echocardiographic dyssynchrony to long-term survival after cardiac resynchronization therapy. *Circulation* 2010;122:1910–8.
- [42] Courtehoux M, Zannad N, Fauchier L, et al. Interventricular delay measurement using equilibrium radionuclide angiography before resynchronization therapy should be performed outside the area of segmental wall motion abnormalities. *Eur J Nucl Med Mol Imaging* 2011;38:239–44.
- [43] Dreger H, Borges AC, Baumann G, et al. Successful reduction of intraventricular asynchrony is associated with superior response to cardiac resynchronization therapy. *Cardiovasc Ultrasound* 2010;8:35.
- [44] Taha N, Zhang J, Ranjan R, et al. Biventricular pacemaker optimization guided by comprehensive echocardiography—preliminary observations regarding the effects on systolic and diastolic ventricular function and third heart sound. *J Am Soc Echocardiogr* 2010;23:857–66.
- [45] Shevchenko OP, Dobrotvorskaia TE, Liakishev AA, et al. Clinico-morphological and echocardiographic parallels in myocardial hypertrophy in patients with ischemic heart disease. *Kardiologiia* 1977;17:37–41.
- [46] Turcott RG, Witteles RM, Wang PJ, et al. Measurement precision in the optimization of cardiac resynchronization therapy. *Circ Heart Fail* 2010;3:395–404.
- [47] Deplagne A, Bordachar P, Reant P, et al. Additional value of three-dimensional echocardiography in patients with cardiac resynchronization therapy. *Arch Cardiovasc Dis* 2009;102:497–508.
- [48] Thomas DE, Yousef ZR, Fraser AG. A critical comparison of echocardiographic measurements used for optimizing cardiac resynchronization therapy: stroke distance is best. *Eur J Heart Fail* 2009;11:779–88.
- [49] Hamaguchi S, Kinugawa S, Tsuchihashi-Makay M, et al. Characteristics, management, and outcomes for patients during hospitalization due to worsening heart failure—a report from the Japanese Cardiac Registry of Heart Failure in Cardiology (JCARE-CARD). *J Cardiol* 2013;62:95–101.
- [50] Strik M, van Middendorp LB, Houthuizen P, et al. Interplay of electrical wavefronts as determinant of the response to cardiac resynchronization therapy in dyssynchronous canine hearts. *Circ Arrhythm Electrophysiol* 2013;6:924–31.
- [51] D'Onofrio A, Botto G, Mantica M, et al. The interventricular conduction time is associated with response to cardiac resynchronization therapy: interventricular electrical delay. *Int J Cardiol* 2013;168:5067–8.
- [52] Dou J, Xia L, Deng D, et al. A study of mechanical optimization strategy for cardiac resynchronization therapy based on an electromechanical model. *Comput Math Methods Med* 2012;948781.
- [53] Celikyurt U, Agacdiken A, Sahin T, et al. Relationship between fragmented QRS and response to cardiac resynchronization therapy. *J Interv Card Electrophysiol* 2012;35:337–42 ([discussion 342]).
- [54] Sperzel J, Brandt R, Hou W, et al. Intraoperative characterization of interventricular mechanical dyssynchrony using electroanatomic mapping system—a feasibility study. *J Interv Card Electrophysiol* 2012;35:189–96.
- [55] Johnson CK, Cberiza SE, Sahar RL, et al. Clinical validation of a real-time data processing system for cardiac output and arterial pressure measurement during intraoperative biventricular pacing optimization. *ASAIO J* 2012;58:191–6.
- [56] Hara H, Oyenuga OA, Tanaka H, et al. The relationship of QRS morphology and mechanical dyssynchrony to long-term outcome following cardiac resynchronization therapy. *Eur Heart J* 2012;33:2680–91.
- [57] Kristiansen HM, Hovstad T, Vollen G, et al. Clinical implication of right ventricular to left ventricular interlead sensed electrical delay in cardiac resynchronization therapy. *Europace* 2012;14:986–93.
- [58] Kristiansen HM, Vollen G, Hovstad T, et al. The impact of left ventricular lead position on left ventricular reverse remodelling and improvement in mechanical dyssynchrony in cardiac resynchronization therapy. *Eur Heart J Cardiovasc Imaging* 2012;13:991–1000.
- [59] Barold SS, Herweg B. Usefulness of the 12-lead electrocardiogram in the follow-up of patients with cardiac resynchronization devices. Part II. *Cardiol J* 2011;18:610–24.
- [60] Golovchiner G, Dorian P, Mangat I, et al. Electrogram-based optimal atrioventricular and interventricular delays of cardiac resynchronization change individually during exercise. *Can J Cardiol* 2011;27:351–7.
- [61] Scalea TM, Hartnett RW, Duncan AO, et al. Central venous oxygen saturation: a useful clinical tool in trauma patients. *J Trauma* 1990;30:1539–43.
- [62] Duncan AM, Lim E, Mebrate Y, et al. Underestimation of duration of ventricular activation by 12-lead ECG compared with direct measurement of activation duration derived from implanted pacemaker leads. *Int J Cardiol* 2011;152:35–42.
- [63] Tamborero D, Mont L, Sitges M, et al. Optimization of the interventricular delay in cardiac resynchronization therapy using the QRS width. *Am J Cardiol* 2009;104:1407–12.
- [64] Muellerleile K, Baholli L, Groth M, et al. Quantification of mechanical ventricular dyssynchrony: direct comparison of velocity-encoded and cine magnetic resonance imaging. *Rofo* 2011;183:554–60.
- [65] Doltra A, Vidal B, Silva E, et al. Comparison of hemodynamic versus dyssynchrony assessment for interventricular delay optimization with echocardiography in cardiac resynchronization therapy. *Pacing Clin Electrophysiol* 2011;34:984–90.
- [66] Kuhne M, Blank R, Schaer B, et al. Effects of physical exercise on cardiac dyssynchrony in patients with impaired left ventricular function. *Europace* 2011;13:839–44.
- [67] Bocchiardo M, Meyer zu Vilsendorf D, Militello C, et al. Resynchronization therapy optimization by intracardiac impedance. *Europace* 2010;12:1589–95.
- [68] Spotnitz ME, Wang DY, Quinn TA, et al. Hemodynamic stability during biventricular pacing after cardiopulmonary bypass. *J Cardiothorac Vasc Anesth* 2011;25:238–42.
- [69] Khan FZ, Virdee MS, Pugh PJ, et al. Non-invasive cardiac output measurements based on bioreactance for optimization of atrio- and interventricular delays. *Europace* 2009;11:1666–74.
- [70] Martinovic SS, Perisic Z, Milic D, et al. Echocardiographic estimation of acute haemodynamic response during optimization of multisite pace-maker using different pacing modalities and atrioventricular delays. *Vojnosanit Pregl* 2009;66:365–70.
- [71] Bogaard MD, Meine M, Tuinenburg AE, et al. Cardiac resynchronization therapy beyond nominal settings: who needs individual programming of the atrioventricular and interventricular delay? *Europace* 2012;14:1746–53.
- [72] Bogaard MD, Meine M, Doevendans PA. Programmed versus effective VV delay during CRT optimization: when what you see is not what you get. *Pacing Clin Electrophysiol* 2013;36:403–9.
- [73] Stegmann B, Francis DP. Atrioventricular and interventricular delay optimization and response quantification in biventricular pacing: arrival of reliable clinical algorithms and research protocols, and how to distinguish them from unreliable counterparts. *Europace* 2012;14:1679–83.
- [74] Lumens J, Leenders GE, Cramer MJ, et al. Mechanistic evaluation of echocardiographic dyssynchrony indices: patient data combined with multiscale computer simulations. *Circ Cardiovasc Imaging* 2012;5:491–9.
- [75] van Deursen CJ, Strik M, Rademakers LM, et al. Vectorcardiography as a tool for easy optimization of cardiac resynchronization therapy in canine left bundle branch block hearts. *Circ Arrhythm Electrophysiol* 2012;5:544–52.
- [76] Cheng CM, Huang JL, Wu TJ, et al. Comparison of quick optimization of interventricular delay between simple methods: intracardiac electrogram and surface electrocardiogram after cardiac resynchronization therapy. *Europace* 2012;14:1317–23.
- [77] Dreger H, Antonow G, Spethmann S, et al. Dyssynchrony parameter-guided interventricular delay programming. *Europace* 2012;14:696–702.
- [78] Bogaard MD, Kirkels JH, HAuer RN, et al. Should we optimize cardiac resynchronization therapy during exercise? *J Cardiovasc Electrophysiol* 2010;21:1307–16.
- [79] Bogaard MD, Doevendans PA, Leenders GE, et al. Can optimization of pacing settings compensate for a non-optimal left ventricular pacing site? *Europace* 2010;12:1262–9.
- [80] Silva E, Sitges M, Mont L, et al. Quantification of left ventricular asynchrony throughout the whole cardiac cycle with a computed algorithm: application for optimizing resynchronization therapy. *J Cardiovasc Electrophysiol* 2009;20:1130–6.
- [81] Miri R, Reumann M, Farina D, et al. Concurrent optimization of timing delays and electrode positioning in biventricular pacing based on a computer heart model assuming 17 left ventricular segments. *Biomed Tech (Berl)* 2009;54:55–65.
- [82] Anselmino M, Antolini M, Amelione C, et al. Optimization of cardiac resynchronization therapy: echocardiographic vs semiautomatic device algorithms. *Congest Heart Fail* 2009;15:14–8.
- [83] Wikstrom G, Blomstrom-Lundqvist C, Andren B, et al. The effects of aetiology on outcome in patients treated with cardiac resynchronization therapy in the CARE-HF trial. *Eur Heart J* 2009;30:782–8.
- [84] Altman M, Edstedt Bonamy AK, Wikstrom AK, et al. Cause-specific infant mortality in a population-based Swedish study of term and post-term births: the contribution of gestational age and birth weight. *BMJ Open* 2012;2 (pii: e001152).
- [85] Andrikopoulos GK, Tzeis S, Kolb C, et al. Correlation of mechanical dyssynchrony with QRS duration measured by signal-averaged electrocardiography. *Ann Noninvasive Electrocardiol* 2009;14:234–41.